

# 1 pKa computation using rMIB

## 1.1 pKa computing procedure

Following the discussion in the previous section, we use the programming language Python to write a wrapper to pipeline the pKa computation in four steps as 1) changing the protonation states on PDB and then on PQR files, with each PQR file representing a needed charge distribution in the pK<sub>a</sub> computation, 2) calling MIBPB solver for solving electrostatics potential and free energies for all PQR files, 3) calculating the intrinsic pK<sub>a</sub> values, and 4) finally titrating the final pK<sub>a</sub> values with energies including site-site interactions. The procedure implemented with the Python wrapper is given as the following.

### Step 1: Prepare the protein structure and protonation states

In this step, we prepare the structure and charge distribution of the protein. The protein structure is obtained from the Protein Data Bank (PDB) ([www.pdb.org](http://www.pdb.org)). The charge distribution is then produced by PDB2PQR [1] with user chosen forcefields. Before calling the PDB2PQR program, the protonation states are configured by using different residue names as specified below in Table 1. For a protein with  $N_t$  titration sites, we will need

Table 1: The protonation states of titratable amino acids: each residue has a name for unprotonated state and a name for protonated states; total charge of the residue in a state is included in the parentheses.

standard resid.	ARG	ASP	GLU	HIS	LYS	TYR	CYS
unprot.(chg)	AR0 (+1)	ASP(-1)	GLU(-1)	HIE(0)	LYN(0)	TYM(-1)	CYM(-1)
protonated(chg)	ARG(+2)	ASH(0)	GLH(0)	HIP(+1)	LYS(+1)	TYR(0)	CYX(0)

$1 + 2N_t + \frac{1}{2}(N_t^2 - N_t)$  charge distributions in the form of PQR files as specified below.

- (1) one PQR file with all titration sites unprotonated, keeping the background charge;
- (2)  $2N_t$  PQR files having all titration sites unprotonated but one protonated with or without the background charge; those with background charges on are used for calculating the intrinsic pK<sub>a</sub>;
- (3)  $\frac{1}{2}(N_t^2 - N_t)$  PQR files with  $i$ th and  $j$ th titration sites protonated only; all background charges are set to zero.

### Step 2: Call MIBPB solver for solving electrostatics

This step calculates the electrostatic free energies for all titration states represented by different PQR files. Each PQR file with different charge distribution changes the RHS of the PB equation. The PB equation can then be accurately and conveniently solved using the MIBPB solver. In calling the MIBPB solver, the user can specify solver and PB model related parameters such as dielectric constants, ion concentration, mesh size, etc. in the *usrdata.in* file. After this step, we received electrostatic free energies from all charge distributions. These energies will be used to calculate pK<sub>a</sub> as explained in the next two steps. Our work follows the procedure as described in Ho's thesis [2].

### Step 3: Compute the intrinsic $pK_a$

The intrinsic  $pK_a$  for the  $i$ th titratable site can be computed by Eq. (1.1) [2],

$$pK_{a,i}^{intr} = pK_{a,i}^0 - \frac{1}{RT \ln 10} [\Delta G_{\text{ele}}(A_p \rightarrow A_p H) - \Delta G_{\text{ele}}(A_s \rightarrow A_s H)] \quad (1.1)$$

where the protein environment has only the fixed background charges, i.e. with all titration sites unprotonated. In this equation,  $\Delta G_{\text{ele}}(A_p \rightarrow A_p H)$  is the difference of the free energy between protein with  $i$ th titration site protonated and protein with all titration unprotonated while  $\Delta G_{\text{ele}}(A_s \rightarrow A_s H)$  is the difference of the free energy between protonated and unprotonated residues alone. Here  $pK_{a,i}^0$  is the model  $pK_a$  at  $T = 298\text{K}$  taken from [3] as in Table 2. For the related constants,  $R$  is the gas constant and  $T$  is the tem-

ARG	ASP	CYS	GLU	HIS	LYS	TYR
12.0	4.0	9.5	4.4	6.3	10.4	9.6

Table 2: model  $pK_a$  value for titration sites.

perature in Kelvin.  $R$  is related to the Boltzmann constant,  $k_B = 1.3806 \cdot 10^{-23} \text{J/K}$ , and the Avogadro constant,  $N_A = 6.02 \cdot 10^{23} / \text{mol}$ , as

$$R = k_B \cdot N_A \approx 8.31 \text{J} / (\text{mol} \cdot \text{K}), \quad (1.2)$$

Since the energy calculated from MIBPB using the unit kcal/mol, we finally use the  $RT$  values as

$$RT \approx 8.31 \cdot 298 \text{J/mol} \approx 2.5 \text{kJ/mol} = (2.5/4.182) \text{kcal/mol}, \quad (1.3)$$

thus the energy from MIBPB divided by  $(2.5 \ln 10 / 4.182 = 1.3765)$  kcal/mol leads to the unit of  $pK_a$  values.

### Step 4: Titrating final $pK_a$ with energies including site-site interactions

Recall that  $pK_a$  of a titration site is defined as the pH value in which half of the site is protonated. In the context of computing  $pK_a$  using electrostatic free energy under different titration states, we are looking for the pH which makes the Boltzmann average  $\langle \theta_i, \text{pH} \rangle$  as in Eq. (1.4) to equal 0.5.

$$\langle \theta_i, \text{pH} \rangle = \frac{\sum_{\theta} \theta_i e^{-\Delta G(A \rightarrow A(\theta); \text{pH})/RT}}{\sum_{\theta} e^{-\Delta G(A \rightarrow A(\theta); \text{pH})/RT}}, \quad (1.4)$$

where  $\theta \in \{0, 1\}^{N_t}$  and  $\theta_i \in \{0, 1\}$  is the  $i$ th entry of  $\theta$ . Note the number of states  $\theta$  increases exponentially ( $2^{N_t}$ ) with number of titrating sites  $N_t$ , thus is computationally prohibitive. In this paper we use direct formulation for proteins with less than 20 titrating sites and we are working on statistical sampling approaches to handle cases with larger  $N_t$  for future work.

In Eq. (1.4), the energy  $\Delta G(A \rightarrow A(\theta); \text{pH})$  is evaluated as

$$\Delta G(A \rightarrow A(\theta); \text{pH}) = -RT \ln 10 \sum_i \theta_i (\text{p}K_{a,i}^{\text{intr}} - \text{pH}) + \frac{1}{2} \sum_i \theta_i \sum_{j \neq i} \theta_j \Delta G_{ij}. \quad (1.5)$$

where  $\text{p}K_{a,i}^{\text{intr}}$  is the intrinsic  $\text{p}K_a$  for the  $i$ th titration site as calculated in step 3 by Eq. (1.1).  $\Delta G_{ij}$  as calculated in Eq. (1.6) is the site-site interaction energy (the free energy of the protein having the  $i$ th site protonated for producing electrostatic potential and the  $j$ th titration site protonated for producing energy without the background charge) computed in step 2 as well.

$$\Delta G_{ij} = t_i^T W t_j = \frac{1}{2} (t_i + t_j)^T W (t_i + t_j) - \frac{1}{2} t_i^T W t_i - \frac{1}{2} t_j^T W t_j \quad (1.6)$$

The second equality holds under the assumption that  $W$  is symmetric.

## 1.2 Software dissemination

With the publication of this manuscript, the python wrapper (*wrapper\_pka.py*) and the binary MIBPB solver (*rMIB.exe* on multiple platforms such as MacOS and Linux/Unix) can be found on the author's website sponsored by Southern Methodist University. Since part of the MIBPB solver source code is copyrighted to Michigan State University, please contact Dr. Guowei Wei if interested. The user also needs the MSMS software [4] for molecular surface generation and we include its binary versions as well. In addition, the users need to install PDB2PQR [1].

To compute  $\text{p}K_a$ , the user specifies PB model and MIBPB related parameters (dielectric constants, ion concentration, mesh size, density of molecular surface triangulation, boundary conditions, method of charge regularization, etc.) in *usrdata.in* file. On a computer with Python and Fortran compilers installed and target protein specified with its four-digit PDB ID, the user runs the program by simply typing:

```
python wrapper_pka.py PDBID
```

The wrapper will download PDB file from the protein data bank, identify all titration sites, call MIBPB solver for electrostatics, and return the computed  $\text{p}K_a$  values.

## References

- [1] T. J. Dolinsky, P. Czodrowski, H. Li, J. E. Nielsen, J. H. Jensen, G. Klebe, and N. A. Baker, "Pdb2pqr: expanding and upgrading automated preparation of biomolecular structures for molecular simulations," *Nucleic Acids Research*, vol. 35, 2007.
- [2] K. L. Ho, *Fast direct methods for molecular electrostatics*. PhD thesis, New York University, 2012.
- [3] Y. Nozaki and C. Tanford, "Examination of titration behavior," *Methods in Enzymology*, vol. 11, pp. 715 – 734, 1967.
- [4] M. F. Sanner, A. J. Olson, and J. C. Spehner, "Reduced surface: An efficient way to compute molecular surfaces," *Biopolymers*, vol. 38, pp. 305–320, 1996.